

IMPROVEMENTS OF TWO PRECONDITIONED AOR ITERATIVE METHODS FOR Z-MATRICES

MOHSEN HASANI AND DAVOD KHOJASTEH SALKUYEH*

Communicated by

ABSTRACT. In (*Linear Algebra Appl.* **93** (1987) 161–170) and (*Trans. Japan Soc. Indust. Appl. Math.* **13** (2003) 439–445), two preconditioned AOR iterative methods have been presented for system of linear equations whose coefficient matrix is a Z-matrix. In this paper, we improve these methods and give some comparison theorems to show that the proposed preconditioners are superior to the primitive ones. Finally some numerical experiments are given to show the effectiveness of the proposed preconditioners.

1. Introduction

Consider the system of linear equations

$$(1.1) \quad Ax = b,$$

where $A = (a_{ij}) \in \mathbb{R}^{n \times n}$ is nonsingular matrix and $b \in \mathbb{R}^n$. A basic iterative method to solve Eq. (1.1) can be written as

$$x^{(k+1)} = M^{-1}Nx^{(k)} + M^{-1}b, \quad k = 0, 1, 2, \dots,$$

in which $A = M - N$, where $M, N \in \mathbb{R}^{n \times n}$ and M is nonsingular. It is well-known that this iterative method converges to the solution of Eq. (1.1) for every initial guess x_0 if and only if $\rho(M^{-1}N) < 1$, where

MSC(2010): Primary: 65F10; Secondary: 65F50.

Keywords: System of linear equations, Preconditioner, AOR iterative method, Z-matrix.

Received: , Accepted: .

*Corresponding author

© 2012 Iranian Mathematical Society.

$\rho(\cdot)$ refers to the spectral radius of matrix. Let $a_{ii} \neq 0$, $i = 1, 2, \dots, n$. Therefore, without loss of generality we can assume that $a_{ii} = 1$, $i = 1, 2, \dots, n$. In this case, we split A into

$$(1.2) \quad A = I - L - U,$$

where I is the identity matrix, $-L$ and $-U$ are strictly lower and strictly upper triangular matrices, respectively. The accelerated overrelaxation (AOR) iterative method to solve Eq. (1.1) is defined by [5]

$$x^{(k+1)} = \mathcal{L}_{r,w}x^{(k)} + w(I - rL)^{-1}b,$$

in which $\mathcal{L}_{r,w} = (I - rL)^{-1}[(1 - w)I + (w - r)L + wU]$, where w and r are real parameters and $w \neq 0$. For certain values of the parameters w and r the AOR iterative method results in the Jacobi, Gauss-Seidel and the SOR methods [5].

To improve the convergence rate of an iterative method one may apply it to the preconditioned linear system $PAx = Pb$, where the matrix P is called a preconditioner. Several preconditioners have been presented for the basic iterative methods by many authors [2-4,6-18]. In this paper, we consider the preconditioners $P_{\alpha,\beta} = I + R_{\alpha,\beta}$ and $P_{\alpha,\beta}^* = I + C_{\alpha,\beta}$, where I is the identity matrix and

$$R_{\alpha,\beta} = \begin{pmatrix} 0 & 0 & \dots & 0 & 0 \\ 0 & 0 & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \dots & 0 & 0 \\ -\alpha a_{n1} + \beta & -\alpha a_{n2} + \beta & \dots & -\alpha a_{nn-1} + \beta & 0 \end{pmatrix},$$

$$C_{\alpha,\beta} = \begin{pmatrix} 0 & 0 & \dots & 0 \\ -\alpha a_{21} + \beta & 0 & \dots & 0 \\ -\alpha a_{31} + \beta & 0 & \dots & 0 \\ \vdots & \vdots & \vdots & \vdots \\ -\alpha a_{n1} + \beta & 0 & \dots & 0 \end{pmatrix},$$

in which α and β are nonnegative real numbers. In [11], Milaszewicz has applied the preconditioner $P_{1,0}^*$ to the Gauss-Seidel method. Then, Morimoto et al. have used the preconditioner $P_{1,0}$ to the Gauss-Seidel method [12]. In this paper, we apply the preconditioners $P_{\alpha,\beta}$ and $P_{\alpha,\beta}^*$ to the AOR iterative method and give some comparison theorems.

For convenience, we first present some notations, definitions and preliminaries which will be used in this paper. A matrix $A = (a_{ij}) \in \mathbb{R}^{n \times n}$

is said to be nonnegative and denoted by $A \geq 0$ if $a_{ij} \geq 0$ for all i and j and A is said to be positive and denoted by $A \gg 0$ if $a_{ij} > 0$ for all i and j .

In the sequel, we recall some definitions, lemmas and theorems for later use.

Theorem 1.1. [1] (*Perron-Frobenius*) *If $A \geq 0$, then $\rho(A)$ is an eigenvalue of A , and corresponding to $\rho(A)$, A has a nonnegative eigenvector (Perron vector).*

Definition 1.2. A matrix $A = (a_{ij}) \in \mathbb{R}^{n \times n}$ is a *Z-matrix* if $a_{ij} \leq 0$ for $i \neq j$.

Definition 1.3. A Z-matrix A is said to be an *M-matrix* if A is nonsingular and $A^{-1} \geq 0$.

Definition 1.4. Let $A \in \mathbb{R}^{n \times n}$. The representation $A = M - N$ is called a *splitting* of A if M is nonsingular. The splitting $A = M - N$ is called

- (a) *convergent* if $\rho(M^{-1}N) < 1$;
- (b) an *M-splitting* of A if M is an M-matrix and $N \geq 0$.

Theorem 1.5. [16, Lemma 1.4] *Let A be a nonnegative matrix.*

- (a) *If $\alpha x \leq Ax$ for some nonnegative vector $x \neq 0$, then $\alpha \leq \rho(A)$.*
- (b) *If $Ax \leq \beta x$ for some positive vector x , then $\rho(A) \leq \beta$. Moreover, if A is irreducible and if $0 \neq \alpha x \leq Ax \leq \beta x$ for some nonnegative vector x , then $\alpha \leq \rho(A) \leq \beta$ and x is positive.*

Lemma 1.6. [16, Lemma 1.5] *Let $A = M - N$ be an M-splitting of A . Then $\rho(M^{-1}N) < 1$ if and only if A is an M-matrix.*

Lemma 1.7. [16, Lemma 1.6] *Let A be a Z-matrix. Then, A is an M-matrix if and only if there is a positive vector x such that $Ax \gg 0$.*

Lemma 1.8. [15, Lemma 2.2] *Let $A = (a_{ij}) \in \mathbb{R}^{n \times n}$ be an M-matrix. Then, there exists $\epsilon_0 > 0$ such that, for any $0 < \epsilon \leq \epsilon_0$, $A(\epsilon) = (a_{ij}(\epsilon))$ is also an M-matrix, where*

$$a_{ij}(\epsilon) = \begin{cases} a_{ij}, & \text{if } a_{ij} \neq 0, \\ -\epsilon, & \text{if } a_{ij} = 0. \end{cases}$$

2. PRECONDITIONED AOR ITERATIVE METHOD WITH $P_{\alpha,\beta}$

For the sake of the simplicity let $R = R_{1,0}$, $R_\alpha = R_{\alpha,0}$, $P = I + R$ and $P_\alpha = I + R_\alpha$. We first consider the preconditioner P_α to the system

(1.1), where α is a nonnegative real number. In this case, the coefficient matrix of the preconditioned system can be written as

$$\bar{A} = (I + R_\alpha)A = I - L - U + R_\alpha - R_\alpha L - R_\alpha U = \bar{D} - \bar{L} - \bar{U},$$

where \bar{D} , \bar{L} and \bar{U} are diagonal, strictly lower triangular and strictly upper triangular matrices, respectively, i.e.,

$$(2.1) \quad \bar{D} = I - E(\alpha), \quad \bar{L} = L - R_\alpha + R_\alpha L + F(\alpha), \quad \bar{U} = U,$$

in which $E(\alpha)$ and $F(\alpha)$ are the diagonal and strictly lower triangular parts of $R_\alpha U$, respectively. If $\bar{D} - r\bar{L}$ is nonsingular, the iteration matrix of the AOR iterative method to solve the preconditioned system $P_\alpha Ax = P_\alpha b$ is of the form

$$\bar{L}_{r,w} = (\bar{D} - r\bar{L})^{-1}[(1-w)\bar{D} + (w-r)\bar{L} + w\bar{U}].$$

Let $\beta \geq 0$. If we apply the preconditioner $P_{\alpha,\beta}$ to the system (1.1), then the coefficient matrix of the system can be written as

$$A' = (I + R_{\alpha,\beta})A = I - L - U + R_{\alpha,\beta} - R_{\alpha,\beta}L - R_{\alpha,\beta}U = D' - L' - U',$$

where D' , L' and U' , are the diagonal, strictly lower triangular and strictly upper triangular matrices, respectively. Assuming $R_\beta = R_{\alpha,\beta} - R_\alpha$, it can be seen that

$$(2.2) \quad D' = \bar{D} - E(\beta), \quad L' = \bar{L} - R_\beta + R_\beta L + F(\beta), \quad U' = \bar{U},$$

where $E(\beta)$ and $F(\beta)$ are the diagonal and strictly lower parts of $R_\beta U$, respectively. Now, if $D' - rL'$ is nonsingular, then the iteration matrix of the AOR iterative method with the preconditioner $P_{\alpha,\beta}$ is of the form

$$L'_{r,w} = (D' - rL')^{-1}[(1-w)D' + (w-r)L' + wU'].$$

Theorem 2.1. *Let A be an M -matrix. If*

$$(1 - \alpha)a_{nj} - \alpha \sum_{\substack{k=1 \\ k \neq j}}^{n-1} a_{nk}a_{kj} + \beta(1 + \sum_{\substack{k=1 \\ k \neq j}}^{n-1} a_{kj}) \leq 0,$$

for $j = 1, \dots, n$, then A' is an M -matrix, too.

Proof. The entries a'_{ij} of $A' = (a'_{ij})$ are given by the expressions

$$a'_{ij} = \begin{cases} a_{ij}, & 1 \leq i < n-1, 1 \leq j \leq n, \\ a_{nj} - \sum_{k=1}^{n-1} (\alpha a_{nk} - \beta) a_{kj}, & i = n, 1 \leq j \leq n. \end{cases}$$

Since A is a Z-matrix, we have $a'_{ij} \leq 0$ for $i \neq n$. Also, for $i = n$ we see that

$$a_{ij} = (1 - \alpha)a_{nj} - \alpha \sum_{\substack{k=1 \\ k \neq j}}^{n-1} a_{nk}a_{kj} + \beta(1 + \sum_{\substack{k=1 \\ k \neq j}}^{n-1} a_{kj}) \leq 0.$$

Therefore, A' is a Z-matrix. By Lemma 1.7 there is a positive vector x such that $Ax \gg 0$. On the other hand, $A'x = (I + R_{\alpha, \beta})Ax \gg 0$. Again, by Lemma 1.7 we conclude that the matrix A' is an M-matrix. \square

Remark 2.2. Let A be an M-matrix. If

$$(1 - \alpha)a_{nj} - \alpha \sum_{\substack{k=1 \\ k \neq j}}^{n-1} a_{nk}a_{kj} \leq 0,$$

for $1 \leq j \leq n$, then \bar{A} is an M-matrix, too.

Proof. It is enough to set $\beta = 0$ in Theorem 2.1. \square

Theorem 2.3. Let $A = (a_{ij}) \in \mathbb{R}^{n \times n}$ be a nonsingular Z-matrix, $0 \leq r \leq w \leq 1$, $w \neq 0$ and

$$(1 - \alpha)a_{nj} - \alpha \sum_{\substack{k=1 \\ k \neq j}}^{n-1} a_{nk}a_{kj} + \beta(1 + \sum_{\substack{k=1 \\ k \neq j}}^{n-1} a_{kj}) \leq 0,$$

for $1 \leq j \leq n$.

(1) If $\rho(L_{r,w}) < 1$, then $\rho(L'_{r,w}) \leq \rho(L_{r,w}) < 1$.

(2) If $\rho(L_{r,w}) > 1$ and $1 - \sum_{k=1}^{n-1} (\alpha a_{nk} - \beta)a_{kn} > 0$, then $\rho(L'_{r,w}) \geq \rho(L_{r,w}) > 1$.

Proof. See Theorem 2.6 in [15]. \square

Remark 2.4. Let $A = (a_{ij}) \in \mathbb{R}^{n \times n}$ be a nonsingular Z-matrix, $0 \leq r \leq w \leq 1$, $w \neq 0$ and

$$(1 - \alpha)a_{nj} - \alpha \sum_{\substack{k=1 \\ k \neq j}}^{n-1} a_{nk}a_{kj} \leq 0,$$

for $1 \leq j \leq n$.

(1) If $\rho(L_{r,w}) < 1$, then $\rho(\bar{L}_{r,w}) \leq \rho(L_{r,w}) < 1$.

(2) If $\rho(L_{r,w}) > 1$ and $1 - \alpha \sum_{k=1}^{n-1} a_{nk}a_{kn} > 0$, then $\rho(\bar{L}_{r,w}) \geq \rho(L_{r,w}) > 1$.

Proof. It is enough to set $\beta = 0$ in Theorem 2.3. \square

In the sequel, we present a comparison theorem concerning the proposed preconditioners.

Theorem 2.5. *Let $A = (a_{ij}) \in \mathbb{R}^{n \times n}$ be a nonsingular Z -matrix, $0 \leq r \leq w \leq 1$, $w \neq 0$ and*

$$(1-\alpha)a_{nj} - \alpha \sum_{\substack{k=1 \\ k \neq j}}^{n-1} a_{nk}a_{kj} + \beta(1 + \sum_{\substack{k=1 \\ k \neq j}}^{n-1} a_{kj}) \leq 0, \quad (1-\alpha)a_{nj} - \alpha \sum_{\substack{k=1 \\ k \neq j}}^{n-1} a_{nk}a_{kj} \leq 0,$$

for $1 \leq j \leq n$.

(1) *If $\rho(L_{r,w}) < 1$, then $\rho(L'_{r,w}) \leq \rho(\bar{L}_{r,w}) < 1$.*

(2) *If $\rho(L_{r,w}) > 1$ and $1 - \sum_{k=1}^{n-1} (\alpha a_{nk} - \beta) a_{kn} > 0$, then $\rho(L'_{r,w}) \geq \rho(\bar{L}_{r,w}) > 1$.*

Proof. Assume that

$$\begin{aligned} A &= M - N = \frac{1}{w}(I - rL) - \frac{1}{w}[(1-w)I + (w-r)L + wU], \\ \bar{A} &= \bar{M} - \bar{N} = \frac{1}{w}(\bar{D} - r\bar{L}) - \frac{1}{w}[(1-w)\bar{D} + (w-r)\bar{L} + w\bar{U}]. \end{aligned}$$

It is easy to see that under the assumptions of the theorem $A = M - N$ is an M-splitting. We consider the two following cases.

Case I: If $\rho(L_{r,w}) < 1$, then by Lemma 1.6, A is an M-matrix, and therefore by Remark 2.2 \bar{A} is an M-matrix as well. Hence, $\bar{D} > 0, \bar{L} \geq 0, \bar{U} \geq 0$ and we see that $\bar{A} = \bar{M} - \bar{N}$ is an M-splitting and as a result by Lemma 1.6 we have $\rho(\bar{L}_{r,w}) < 1$. On the other hand, we have

$$\begin{aligned} \bar{L}_{r,w} &= (\bar{D} - r\bar{L})^{-1}[(1-w)\bar{D} + (w-r)\bar{L} + w\bar{U}] \\ &= (I - r\bar{D}^{-1}\bar{L})^{-1}[(1-w)I + (w-r)\bar{D}^{-1}\bar{L} + w\bar{D}^{-1}\bar{U}] \\ &= [I + (r\bar{D}^{-1}\bar{L}) + (r\bar{D}^{-1}\bar{L})^2 + \dots] \\ &\quad \times [(1-w)I + (w-r)\bar{D}^{-1}\bar{L} + w\bar{D}^{-1}\bar{U}] \\ (2.3) \quad &\geq [(1-w)I + (w-r)\bar{D}^{-1}\bar{L} + w\bar{D}^{-1}\bar{U}], \end{aligned}$$

which shows that the matrix $\bar{L}_{r,w}$ is nonnegative. By Theorem 1.1 there is a nonnegative vector x such that $\bar{L}_{r,w}x = \rho(\bar{L}_{r,w})x$. For the sake of the simplicity, let $\lambda = \rho(\bar{L}_{r,w})$. Then

$$[(1-w)\bar{D} + (w-r)\bar{L} + w\bar{U}]x = \lambda(\bar{D} - r\bar{L})x,$$

and

$$(2.4) \quad wU'x = w\bar{U}x = (\lambda - 1 + w)\bar{D}x + (r - w - \lambda r)\bar{L}x.$$

We also have

$$\begin{aligned} \lambda(D' - rL')x &= \lambda(1-r)D'x + \lambda r(D' - L')x \\ (2.5) \quad &= \lambda(1-r)D'x + \lambda r[\bar{D} - E(\beta) - \bar{L} + R_\beta - R_\beta L - F(\beta)]x. \end{aligned}$$

By using Eqs. (2.1), (2.2), (2.4) and (2.5), we have

$$\begin{aligned} L'_{r,w}x - \lambda x &= (D' - rL')^{-1}[(1-w)D' + (w-r)L' + wU' - \lambda(D' - rL')]x \\ &= (D' - rL')^{-1}[(1-w)D' + (w-r)L' + (\lambda-1+w)\bar{D} \\ &\quad + (r-w-\lambda r)\bar{L} - \lambda(1-r)D' - \lambda r(\bar{D} - E(\beta) - \bar{L} \\ &\quad + R_\beta - R_\beta L - F(\beta))]x \\ &= (D' - rL')^{-1}[(1-w-\lambda+\lambda r)D' + \lambda(\bar{D} - r\bar{L}) - (1-w)\bar{D} \\ &\quad + (w-r)(L' - \bar{L}) - \lambda r(\bar{D} - E(\beta) - \bar{L} + R_\beta - R_\beta L - F(\beta))]x \\ &= (D' - rL')^{-1}[(1-w-\lambda+\lambda r)D' + \lambda(\bar{D} - r\bar{L}) - (1-w)\bar{D} \\ &\quad + (w-r)(D' - \bar{D} + E(\beta) - R_\beta + R_\beta L + F(\beta)) - \lambda r(\bar{D} - E(\beta) \\ &\quad - \bar{L} + R_\beta - R_\beta L - F(\beta))]x \\ &= (D' - rL')^{-1}[(\lambda-1)(1-r)(\bar{D} - D') + (w-r+\lambda r)(E(\beta) - R_\beta \\ &\quad + R_\beta L + F(\beta))]x \\ &= (D' - rL')^{-1}[(\lambda-1)(1-r)(\bar{D} - D') \\ &\quad + (w-r+\lambda r)(R_\beta U + R_\beta L - R_\beta)]x \\ &= (D' - rL')^{-1}[(\lambda-1)(1-r)(\bar{D} - D') + (r-\lambda r)R_\beta \\ &\quad + (\lambda r-r)(R_\beta U + R_\beta L) - wR_\beta + wR_\beta L + wR_\beta \bar{U}]x \\ &= (D' - rL')^{-1}[(\lambda-1)(1-r)(\bar{D} - D') + (r-\lambda r)R_\beta \\ &\quad + (\lambda r-r)(R_\beta U + R_\beta L) - wR_\beta + wR_\beta L \\ &\quad + R_\beta((\lambda-1+w)\bar{D} + (r-w-\lambda r)\bar{L})]x \\ &= (D' - rL')^{-1}[(\lambda-1)(1-r)E(\beta) - r(\lambda-1)R_\beta + r(\lambda-1)R_\beta U \\ &\quad + (\lambda-1)R_\beta \bar{D} + wR_\beta \bar{D} - wR_\beta + (w-r+\lambda r)R_\beta(L - \bar{L})]x. \end{aligned}$$

Since $R_\beta E(\alpha) = R_\beta F(\alpha) = R_\beta R_\alpha = 0$, $\bar{D} = I - E(\alpha)$ and $L - \bar{L} = R_\alpha - R_\alpha L - F(\alpha)$, we see that

$$(2.6) \quad L'_{r,w}x - \lambda x = (\lambda-1)(D' - rL')^{-1}[(1-r)E(\beta) + (1-r)R_\beta + rR_\beta U].$$

Let $B = (1 - r)E(\beta) + (1 - r)R_\beta + rR_\beta U$. Obviously, $B \geq 0$. On the other hand, by Theorem 2.1 A' is also an M-matrix and therefore

$$(D' - rL')^{-1} \geq 0.$$

Now, if A is an irreducible matrix, then by Eq. (2.3) we have

$$\bar{L}_{r,w} \geq (1 - w)I + w(1 - r)\bar{D}^{-1}L + w\bar{D}^{-1}U.$$

This relation shows that if $0 \leq r < 1$ (note that $w \neq 0$), then $\bar{L}_{r,w}$ is also irreducible. Hence, by Lemma 4 in [15] we have $\lambda > 0$ and the Perron vector x is positive. Now, from Eq. (2.6) we conclude that

$$L'_{r,w}x \leq \lambda x,$$

and by using Theorem 1.5 we get

$$\rho(L'_{r,w}) \leq \lambda = \rho(\bar{L}_{r,w}).$$

If $r = 1$, then $w = r = 1$ and

$$\rho(L'_{1,1}) = \lim_{r \rightarrow 1^-} \rho(L'_{r,1}) \leq \lim_{r \rightarrow 1^-} \rho(\bar{L}_{r,1}) = \rho(\bar{L}_{1,1}) < 1.$$

On the other hand, if A is a reducible matrix, then by Lemma 1.8 for every small enough $\epsilon > 0$ the matrix $A(\epsilon)$ is an irreducible M-matrix. Hence, we have

$$\rho(L'_{r,w}) = \lim_{\epsilon \rightarrow 0^+} \rho(L'_{r,w}(\epsilon)) \leq \lim_{\epsilon \rightarrow 0^+} \rho(\bar{L}_{r,w}(\epsilon)) = \rho(\bar{L}_{r,w}) < 1.$$

This completes the first part of the theorem.

Case II: Let $\rho(L_{r,w}) > 1$. By Remark (2.4), we have $\rho(\bar{L}_{r,w}) \geq \rho(L_{r,w}) > 1$. Then by the assumption $1 - \sum_{k=1}^{n-1} (\alpha a_{nk} - \beta) a_{kn} > 0$ of the theorem we see that A' is a Z-matrix and therefore $D' - rL'$ is an M-matrix and by Eq. (2.6) and Theorem 1.5, we get

$$\rho(L'_{r,w}) \geq \rho(\bar{L}_{r,w}) > 1.$$

This proves the second part of the theorem. \square

3. PRECONDITIONED AOR ITERATIVE METHOD WITH $P_{\alpha,\beta}^*$

For simplicity, we assume that $C = C_{1,0}$, $C_\alpha = C_{\alpha,0}$, $P^* = I + C$ and $P_\alpha^* = I + C_\alpha$. As we mentioned Milaszewicz in [11] applied preconditioned Gauss-Seidel method in conjunction with the preconditioner P^*

to solve Eq. (1.1). We first consider the preconditioner P_α^* . In this case, $\tilde{A} = P_\alpha^*A$ can be written as

$$\tilde{A} = I - L - U + C_\alpha - C_\alpha U = \tilde{D} - \tilde{L} - \tilde{U},$$

where \tilde{D} , \tilde{L} and \tilde{U} are, respectively, diagonal, strictly lower triangular and strictly upper triangular matrices and

$$(3.1) \quad \tilde{D} = I - E_2(\alpha), \quad \tilde{L} = L - C_\alpha + F_2(\alpha), \quad \tilde{U} = U + G_2(\alpha),$$

in which $E_2(\alpha)$, $F_2(\alpha)$ and $G_2(\alpha)$ are, respectively, diagonal, strictly lower triangular and strictly upper triangular matrices such that

$$C_\alpha U = G_2(\alpha) + E_2(\alpha) + F_2(\alpha).$$

If $\tilde{D} - r\tilde{L}$ is nonsingular, then the iteration matrix of the AOR method with the preconditioner P_α^* can be expressed as

$$\tilde{L}_{r,w} = (\tilde{D} - r\tilde{L})^{-1}[(1-w)\tilde{D} + (w-r)\tilde{L} + w\tilde{U}].$$

Now, we use the preconditioner $P_{\alpha,\beta}^*$. In this case, $\hat{A} = P_{\alpha,\beta}^*A$ can be decomposed as

$$\hat{A} = I - L - U + C_{\alpha,\beta} - C_{\alpha,\beta}U = \hat{D} - \hat{L} - \hat{U},$$

where \hat{D} , \hat{L} and \hat{U} are, respectively, diagonal, strictly lower triangular and strictly upper triangular matrices. If we assume $C_\beta = C_{\alpha,\beta} - C_\alpha$, then

$$(3.2) \quad \hat{D} = \tilde{D} - E_2(\beta), \quad \hat{L} = \tilde{L} - C_\beta + F_2(\beta), \quad \hat{U} = \tilde{U} + G_2(\beta),$$

in which $E_2(\beta)$, $F_2(\beta)$ and $G_2(\beta)$ are, respectively, diagonal, strictly lower triangular and strictly upper triangular matrices such that

$$C_\beta U = G_2(\beta) + E_2(\beta) + F_2(\beta).$$

If $(\hat{D} - r\hat{L})$ is nonsingular, then the iteration matrix of the AOR iterative method with the preconditioner $P_{\alpha,\beta}^*$ can be written as

$$\hat{L}_{r,w} = (\hat{D} - r\hat{L})^{-1}[(1-w)\hat{D} + (w-r)\hat{L} + w\hat{U}].$$

Theorem 3.1. *Let A be an M -matrix. If*

$$(1 - \alpha)a_{i1} + \beta \leq 0,$$

for $2 \leq i \leq n$, then \hat{A} is an M -matrix, too.

Proof. Let A be an M-matrix and $\widehat{A} = (\widehat{a}_{ij})$. Then, we have

$$\widehat{a}_{ij} = a_{ij} + (-\alpha a_{i1} + \beta)a_{1j} = a_{ij} - \alpha a_{i1}a_{1j} + \beta a_{1j},$$

for $2 \leq i \leq n$. Since A is a Z-matrix, obviously we have $\widehat{a}_{ij} \leq 0$, for $i \neq j$ and $j \neq 1$. For $j = 1$, we see that

$$\widehat{a}_{ij} = (1 - \alpha)a_{i1} + \beta \leq 0.$$

Hence, \widehat{A} is a Z-matrix. By Lemma 1.7, there exists a positive vector x such that $Ax \gg 0$. On the other hand, we have $\widehat{A}x = (I + C_{\alpha, \beta})Ax \gg 0$. Therefore, by Lemma 1.7, \widehat{A} is also an M-matrix. \square

Remark 3.2. Let A be an M-matrix. If

$$(1 - \alpha)a_{i1} \leq 0,$$

for $2 \leq i \leq n$, then \widetilde{A} is an M-matrix, too.

Proof. It is enough to set $\beta = 0$ in Theorem 3.1. \square

Theorem 3.3. Let $A = (a_{ij}) \in \mathbb{R}^{n \times n}$ be a nonsingular Z-matrix, $0 \leq r \leq w \leq 1$, $w \neq 0$ and

$$(1 - \alpha)a_{i1} + \beta \leq 0,$$

for $2 \leq i \leq n$.

(1) If $\rho(L_{r,w}) < 1$, then $\rho(\widehat{L}_{r,w}) \leq \rho(L_{r,w}) < 1$.

(2) If $\rho(L_{r,w}) > 1$ and $1 - \alpha a_{i1}a_{1i} + \beta a_{1i} > 0$, then $\rho(\widehat{L}_{r,w}) \geq \rho(L_{r,w}) > 1$.

Proof. See Theorem 2.6 in [15]. \square

Remark 3.4. Let $A = (a_{ij}) \in \mathbb{R}^{n \times n}$ be a nonsingular Z-matrix, $0 \leq r \leq w \leq 1$, $w \neq 0$ and

$$(1 - \alpha)a_{i1} \leq 0,$$

for $2 \leq i \leq n$.

(1) If $\rho(L_{r,w}) < 1$, then $\rho(\widetilde{L}_{r,w}) \leq \rho(L_{r,w}) < 1$.

(2) If $\rho(L_{r,w}) > 1$ and $1 - \alpha a_{i1}a_{1i} > 0$, then $\rho(\widetilde{L}_{r,w}) \geq \rho(L_{r,w}) > 1$.

Proof. It is enough to set $\beta = 0$ in Theorem 3.3. \square

In continuation, we present a comparison theorem concerning the proposed preconditioners.

Theorem 3.5. Let $A = (a_{ij}) \in \mathbb{R}^{n \times n}$ be a nonsingular Z-matrix, $0 \leq r \leq w \leq 1$, $w \neq 0$ and

$$(1 - \alpha)a_{i1} + \beta \leq 0,$$

for $2 \leq i \leq n$.

(1) If $\rho(L_{r,w}) < 1$, then $\rho(\widehat{L}_{r,w}) \leq \rho(\widetilde{L}_{r,w}) < 1$.

(2) If $\rho(L_{r,w}) > 1$ and $1 - \alpha a_{i1} a_{1i} + \beta a_{1i} > 0$, then $\rho(\widehat{L}_{r,w}) \geq \rho(\widetilde{L}_{r,w}) > 1$.

Proof. The proof is quite similar to that of Theorem 2.5 and is omitted here. \square

4. NUMERICAL EXAMPLES

In this section, to evaluate the efficiency of the preconditioners presented in this paper we compare the spectral radii of the iteration matrix of preconditioned AOR iterative methods in conjunction with the proposed preconditioners for two matrices.

Example 4.1. In this example, we consider

$$A = \begin{pmatrix} 1 & -0.1 & -0.1 & -0.1 \\ -0.5 & 1 & 0 & -0.1 \\ -0.5 & 0 & 1 & -0.4 \\ -0.7 & -0.2 & -0.4 & 1 \end{pmatrix},$$

as the coefficient matrix of the system (1.1). For $w = 0.8$, $\alpha = 0.6$, $\beta = 0.1$ and $r = 0.1, 0.2, \dots, 0.9$, it is easy to verify that all the assumptions of the Theorems 2.5 and 3.5 are satisfied. Hence, we expect that $P_{\alpha,\beta}$ (resp. $P_{\alpha,\beta}^*$) is superior to P_α (resp. P_α^*) which is itself better than the preconditioner $P = I$. To confirm this claim we report the spectral radii of the iteration matrix of the preconditioned AOR methods with the proposed preconditioners in TABLE 1. As we observe, the results confirm the claim. For more investigation, let $w = 0.7$, $\alpha = 0.3$, $\beta = 0.15$ and $r = 0.1, 0.2, \dots, 0.9$. We report the results in TABLE 2. As we see, the results give the same conclusion as TABLE 1.

Example 4.2. Let

$$A = \begin{pmatrix} 1 & -0.5 & -0.2 & 0 & -1 \\ -0.2 & 1 & 0 & -0.2 & -0.2 \\ -0.2 & -0.2 & 1 & -0.2 & -0.3 \\ -0.4 & 0 & -0.1 & 1 & -0.1 \\ -0.3 & -0.1 & -0.1 & -0.1 & 1 \end{pmatrix}.$$

Similar to the previous example we present the spectral radii of the iteration matrix of the preconditioned AOR methods with the proposed

TABLE 1. Comparison of spectral radii of the AOR method with preconditioners $P = I, P_\alpha, P_{\alpha,\beta}, P_\alpha^*, P_{\alpha,\beta}^*$ with $w = 0.8$, $\alpha = 0.6$ and $\beta = 0.1$ for different values of r for Example 4.1.

r	$P = I$	$P = P_\alpha$	$P = P_{\alpha,\beta}$	$P = P_\alpha^*$	$P = P_{\alpha,\beta}^*$
0.1	0.6873	0.6125	0.5697	0.6375	0.6188
0.2	0.6760	0.6004	0.5579	0.6255	0.6066
0.3	0.6636	0.5871	0.5450	0.6123	0.5933
0.4	0.6498	0.5725	0.5309	0.5977	0.5787
0.5	0.6344	0.5563	0.5152	0.5815	0.5623
0.6	0.6169	0.5379	0.4976	0.5631	0.5438
0.7	0.5968	0.5167	0.4773	0.5419	0.5224
0.8	0.5731	0.4918	0.4535	0.5167	0.4971
0.9	0.5440	0.4609	0.4240	0.4856	0.4656

TABLE 2. Comparison of spectral radii of the AOR method with preconditioners $P = I, P_\alpha, P_{\alpha,\beta}, P_\alpha^*, P_{\alpha,\beta}^*$ with $w = 0.7$, $\alpha = 0.3$ and $\beta = 0.15$ for different values of r for Example 4.1.

r	$P = I$	$P = P_\alpha$	$P = P_{\alpha,\beta}$	$P = P_\alpha^*$	$P = P_{\alpha,\beta}^*$
0.1	0.7568	0.7311	0.6909	0.7393	0.7214
0.2	0.7480	0.7219	0.6818	0.7303	0.7122
0.3	0.7383	0.7119	0.6719	0.7203	0.7020
0.4	0.7276	0.7008	0.6610	0.7093	0.6909
0.5	0.7156	0.6884	0.6488	0.6970	0.6784
0.6	0.7021	0.6744	0.6352	0.6830	0.6643
0.7	0.6864	0.6583	0.6195	0.6669	0.6480
0.8	0.6679	0.6392	0.6011	0.6480	0.6288
0.9	0.6453	0.6158	0.5784	0.6246	0.6051

preconditioners in TABLES 3 and 4 for two set of parameters. In the first set we consider $w = 0.9$, $\alpha = 0.3$, $\beta = 0.1$ and $r = 0.1, 0.2, \dots, 0.9$ and for the second set $w = 0.9$, $\alpha = 0.4$, $\beta = 0.12$ and $r = 0.1, 0.2, \dots, 0.9$. Both of TABLES 3 and 4 show that the preconditioner $P_{\alpha,\beta}$ (resp. $P_{\alpha,\beta}^*$) is better than P_α (resp. P_α^*) which is itself better than the preconditioner $P = I$.

TABLE 3. Comparison of spectral radii of the AOR method with preconditioners $P = I, P_\alpha, P_{\alpha,\beta}, P_\alpha^*, P_{\alpha,\beta}^*$ with $w = 0.9$, $\alpha = 0.3$ and $\beta = 0.1$ for different values of r for Example 4.1.

r	$P = I$	$P = P_\alpha$	$P = P_{\alpha,\beta}$	$P = P_\alpha^*$	$P = P_{\alpha,\beta}^*$
0.1	0.8971	0.8846	0.8523	0.8818	0.8554
0.2	0.8920	0.8789	0.8457	0.8761	0.8487
0.3	0.8864	0.8727	0.8385	0.8698	0.8414
0.4	0.8801	0.8657	0.8304	0.8627	0.8332
0.5	0.8730	0.8578	0.8213	0.8547	0.8240
0.6	0.8649	0.8488	0.8111	0.8456	0.8135
0.7	0.8556	0.8384	0.7993	0.8352	0.8015
0.8	0.8447	0.8262	0.7856	0.8230	0.7875
0.9	0.8317	0.8116	0.7694	0.8084	0.7708

TABLE 4. Comparison of spectral radii of the AOR method with preconditioners $P = I, P_\alpha, P_{\alpha,\beta}, P_\alpha^*, P_{\alpha,\beta}^*$ with $w = 0.9$, $\alpha = 0.4$ and $\beta = 0.12$ for different values of r for Example 4.1.

r	$P = I$	$P = P_\alpha$	$P = P_{\alpha,\beta}$	$P = P_\alpha^*$	$P = P_{\alpha,\beta}^*$
0.1	0.8971	0.8797	0.8342	0.8756	0.8376
0.2	0.8920	0.8738	0.8271	0.8696	0.8304
0.3	0.8864	0.8673	0.8193	0.8630	0.8224
0.4	0.8801	0.8601	0.8106	0.8556	0.8135
0.5	0.8730	0.8518	0.8009	0.8473	0.8035
0.6	0.8649	0.8425	0.7900	0.8378	0.7921
0.7	0.8556	0.8317	0.7774	0.8269	0.7791
0.8	0.8447	0.8190	0.7628	0.8142	0.7640
0.9	0.8317	0.8039	0.7456	0.7990	0.7459

5. CONCLUSION

We have improved two known preconditioned AOR iterative methods. Some comparison theorems have been given to show the improvement of the preconditioners. Some numerical examples have been presented to validate the presented theoretical results. Numerical results show that

the proposed preconditioners are more effective than the primary version of them.

Acknowledgments

The authors would like to thank the anonymous referee for helpful comments and suggestions.

REFERENCES

- [1] A. Berman and R. J. Plemmons, *Nonnegative Matrices in the Mathematical Sciences*, SIAM, Philadelphia, PA, 1994.
- [2] M. Dehghan and M. Hajarian, Improving preconditioned SOR-type iterative methods for L-matrices, *Commun. Numer. Methods Eng.* **27** (2011) 774–784.
- [3] D. J. Evans, M. M. Martins and M. E. Trigo, The AOR iterative method for new preconditioned linear systems, *J. Comput. Appl. Math.* **132** (2001) 461–466.
- [4] A. D. Gunawardena, S. K. Jain and L. Snyder, Modified iterative methods for consistent linear systems, *Linear Algebra Appl.* **154-156** (1991) 123–143.
- [5] A. Hadjidimos, Accelerated overrelaxation method, *Mathematics of Computation* **32** (1978) 149–157.
- [6] A. Hadjidimos, D. Noutsos and M. Tzoumas, More on modifications and improvements of classical iterative schemes for M-matrices, *Linear Algebra Appl.* **364** (2003) 253–279.
- [7] T. Kohno and H. Kotakemori, Improving the modified Gauss-Seidel method for Z-matrices, *Linear Algebra Appl.* **267** (1997) 113–123.
- [8] H. Kotakemori, H. Niki and N. Okamoto, Accelerated iterative method for Z-matrices, *J. Comput. Appl. Math.* **75** (1996) 87–97.
- [9] W. Li, A note on the preconditioned Gauss-Seidel (GS) method for Linear systems, *J. Comput. Appl. Math.* **182**(2005) 81–90.
- [10] W. Li and W. Sun, Modified Gauss-Seidel type methods and Jacobi type methods for Z-matrices, *Linear Algebra Appl.* **317** (2000) 227–240.
- [11] J. P. Milaszewicz, Improving Jacobi and Gauss-Seidel iterations, *Linear Algebra Appl.* **93** (1987) 161–170.
- [12] M. Morimoto, H. Kotakemori, T. Kohno and H. Niki, The Gauss-Seidel method with preconditioner $(I + R)$, *Trans. Japan Soc. Indust. Appl. Math.* **13** (2003) 439–445.
- [13] L. Wang, On a class of row preconditioners for solving Linear systems, *Inter. J. Comput. Math.* **83** (2006) 939–949.
- [14] H. Wang and Y. T. Li, A new Preconditioned AOR iterative method for L-matrices, *J. Comput. Appl. Math.* **229** (2009) 47–53.
- [15] L. Wang and Y.Z. Song, Preconditioned AOR iterative method for M-matrices, *J. Comput. Appl. Math.* **226** (2009) 114–124.

- [16] M. Wu, L. Wang and Y. Song, Preconditioned AOR iterative method for linear systems *Appl. Numer. Math.* **57** (2007) 672–685.
- [17] J. H. Yun, Comparison results of the Preconditioned AOR methods for L-matrices, *Appl. Math. Comput.* **218** (2011) 3399–3413.
- [18] Y. Zhang, T.Z. Huang, X. P.Liu, Modified iterative methods for nonnegative matrices and M-matrices linear systems, *Comput. Math. Appl.* **50** (2005) 1587–1602.

Mohsen Hasani

Faculty of Science, Department of Mathematics, Islamic Azad University, Shahrood, Iran

Email: hasani.mo@gmail.com

Davod Khojasteh Salkuyeh

Faculty of Mathematical Sciences, University of Guilan, P.O. Box. 1914, Rasht, Iran

Email: khojasteh@guilan.ac.ir, salkuyeh@gmail.com